



REVIEW ARTICLE

Status of research results in chemistry of biologically active substances in Benin

Alain Y. Koudoro, Christian T. R. Konfo, Philippe Sessou and Dominique C. K. Sohounhloué*

Abstract

Research on biologically active chemicals extracted from medicinal plants and essential oils from aromatic plants in the West African subregion is booming. Recognition of the clinical, pharmaceutical and economic value of herbal medicines continues to grow despite the growth of the pharmaceutical industry and the continued development of new, more effective synthetic and biological medical products. On the other hand, despite the improvement in food preservation technics, food preservatives nature remains one of the most important issues for public health. Indeed, several synthetic preservatives have been banned in some countries because of their long-term adverse toxicological effects. The current trend of consumers to seek for a more natural diet has prompted the research, development and application of new natural products with antimicrobial and antioxidant activities in order to use them as alternatives to synthetic preservatives. This review aims to do an inventory of the results of research in chemistry of biologically active substances in Benin.

Keywords: Biologically active substances; food preservation; human and veterinary; pharmacopoeias; plant extracts; public health

Introduction

Medicinal plants are a valuable heritage for humanity, and especially for the majority of poor communities in developing countries who rely on their primary health care and their subsistence [1]. The last twenty years have been seeing an increasing interest in both developed and developing countries for medicinal products [2]. Africa has a significant diversity of medicinal plants. Indeed, on the 300,000 species of species found on the planet more than 200,000 species are found in tropical countries in Africa and have medicinal properties [3]. In Benin, about 3,000 species invented in ecosystems [4], 172 are consumed by local populations such as food plants [5] and 814 as medicinal plants. In fact, the use of traditional medicine is widespread and

has a significant health and economic significance in the country [6]. Chemistry and biology play a central role in the study and valorization of these natural resources. Chemistry, as a discipline of experimental sciences by its structural character, federator and integrator, allows extracting, characterizing and providing health-related substances. Biological applications are, however, essential for a good recovery of these resources. In Benin, several studies focused on the valorization of medicinal plants. These studies were based on ethnobotanical and ethnopharmacological surveys, determining the chemical composition of the volatile and non-volatile extracts of the various plant organs as well as the evaluation of their biological activities such as anti-inflammatory, antibiotics, antimicrobial, anticancer activities and their use as natural food conservatives [7 - 10]. Generally, each plant material studied has been collected and identified with the help of botanists and registered under a number at the National Herbarium of the University of Abomey-Calavi. This review aims to identify biologically active chemicals research results in Benin. It first deals with the biological properties of

* Correspondence: dominique.sohounhloue@uac.bj

Ecole Polytechnique d'Abomey-Calavi, Laboratoire d'Etude et de Recherche en Chimie Appliquée, Université d'Abomey-Calavi, 01 BP 2009, Cotonou, Bénin

Full list of author information is available at the end of the article.

Received: 02 Apr 2018, Accepted: 03 Jun 2018



plants used in human, veterinary pharmacopoeia and their use in the fight against post-harvest deprecators, and in the second time potential food applications of these plant species.

Biological Properties of Plants Used in Pharmacopoeia Human, Veterinary and Pest Control in Post-Harvest in Benin

Use of plants in human pharmacopoeia

In Benin, the most frequent diseases or diseases that cause most morbidity problems, temporary stopping work and mortality are many (malaria, diarrhea, broncho-pulmonary affections, infected traumatic wounds, malnutrition, cardiovascular disease, diabetes and cancers ...). To deal with these problems, medicinal plants are required. Thus *Hyptis suaveolens* are mentioned among the plants used for treating malaria while *Eucalyptus citriodora* is sought to solve the problems of cardiovascular disease, diabetics, obesity and cough etc. Other parts, extracts from *Ceiba pentandra* and *Bridellia ferruginea* are required to treat diarrhea. Other studies [11 - 12] have taken into account the evaluation of the antimicrobial properties of *Lippia rugosa* essential oils and aqueous and ethanolic extracts of *Sebastiania chamaelea*, a plant used by traditional healers in south-central Niger to treat malaria and to relieve dental flare-ups in infants, against reference bacterial strains *Escherichia coli* ATCC 25922 and *Staphylococcus aureus* ATCC 25923 on the one hand and the antifungal properties of *Sebastiania chamaelea* against *Candida albicans* ATCC 10231 on the other hand. The phytochemical screening of *Sebastiania chamaelea* revealed the presence of tanins, polyphenols, flavonoids, triterpens, steroids and traces of quinonic compounds. As for the antibacterial activity of the *Lippia* essential oils, concentrations ranged from 0.44 to 0.88 $\mu\text{L} / \text{mL}$ for the two infected bacteria. In addition, the ethanolic extract of *Chamaelea Sebastiania* has been more active on the studied strains than the aqueous extract which has shown very low antimicrobial activity. *E. coli* and *C. albicans* strains showed low-moderate sensitivity with ethanolic extract (9.3 ± 1.5 mm and 11.33 ± 0.55 mm respectively). *S. aureus*, on the other hand, showed a very significant sensibility (14 ± 1.7 mm inhibition diameter) with ethanolic extract with a Minimal Inhibitory Concentration (MIC) of 3.12 mg.mL^{-1} and a Minimum Bactericidal Concentration (MBC) of 12.50 mg.mL^{-1} . It should be noted that extracts from *Hemizygia bracteosa* were also mentioned among the plants used to combat diabetes in Benin [13].

Use of plants in veterinary pharmacopoeia

The use of plants in the treatment of pathologies is a common practice among pastoralists in Benin. In 2016, a study identified the formulations used by farmers to treat foot-and-mouth disease. The survey was conducted by 370 breeders in diverse

departments in Benin, between July and December 2015 and revealed to treat foot-and-mouth disease, almost all the surveyed breeders associate herbal formulas with veterinary care. A total of 32 types of formulas using 32 plant species have been summarized among breeders. *Vitellaria paradoxa* is the most quoted plant species in 3 types of formulas. Plant species such as *Citrus limon* L., *Gossypium arboreum* L., *Pterocarpus erinaceus*, *Sorghum bicolor* L., *Acacia nilotica* L., *Lannea acida*, *Khaya senegalensis*, *Agelanthus dodoneifolius* were involved in two formulations. The shrubs were involved in 14 formulations and the maceration was the most used method with 27%, followed by the powder (24%) [14].

Another study identified veterinary formulations and selected the most credible before assessing the vulnerability of the plants involved. Most represented families in cash were: leguminosae (34%) and combretaceae (11%). In total 86 recipes were reported to treat 31 pathologies with the most commonly used Pest (19%). Formulations frequencies and quotes allowed to select those that are more credible. The stalks of rod were more cited (61%). Eight (08) methods of preparation were indicated with decoction (49%) as the most frequent. Thirty and one (31) species have been found vulnerable [15].

Another study has listed a range of medicinal plants commonly used to treat animal parasitic pathologies. For this purpose, an ethno-botanical survey was conducted with 787 breeders and agro-farmers of the eight agro-ecological zones of Benin. In total, five (5) medicinal plants (*Azadirachta indica*, *Cajanus cajan*, *Citrus quadrangularis*, *Nicotiana tabacum* and *Vernonia amygdalina*) are used to treat coccidiosis against two (2) plants (*Bombax costatum* and *Afsana prosopis*) for piroplasmose and 12 (12) plants, including *A. polyacantha*, *C. sieberiana*, *C. febrifuga*, *D. microcarpum*, *P. kotschy*, *Z. mays* for trypanosomiasis. In addition, eighteen (18) plants, including *A. digitata*, *A. senegalensis*, *A. leiocarpa*, *Bambusa vulgaris*, *F. exasperata* were invented to treat ectoparasitoses against seventeen (17) plants including *A. flagellaris*, *C. papaya*, *E. guineensis*, *M. charantia*, *Senna alata* for forty and two (42) plants including *A. digitata*, *A. sativum*, *Anacardium occidentale*, *A. leiocarpa*, *B. ferruginea*, *C. papaya*, *C. ambrosioids*, *K. senegalensis* on, *M. inermis* and *N. canensis* to treat helminthiases. The most used organs were sheets and shells. The majority of remedies were prepared as decoration [16].

Finally, extracts from species such as *Ximenia americana*, *Khaya senegalensis* and *Daniellia oliveri* are used to treat gastrointestinal diseases of animals. The African Porcine Pest is treated by the root of *Cochlospermum planchonii* while salmonellosis is treated with *Cissus quadrangularis*. Extracts from *Pseudocedrales kotschy* and *Khaya senegalensis* were

used in the traditional medicine of trypanosomiasis in livestock breeding animals [17].

Plant species used in the fight against pests in food post harvest

Benin's problems by producers during the post-harvest phase of agricultural products have long been negligible with those related to production. Meanwhile, post-harvest losses are becoming increasingly growing, since traditional technology storage and processing technologies are generally inaccurate with risks of infestation of stored products [18]. Chemical control has always been the main measure to reduce the impact of contamination after harvesting. However, applying to high doses of these synthetic chemicals in a diet storage control perspective increases the risk of presence of toxic residues in these products.

The use of plants in crop conservation has been practiced even before the appearance of synthetic insecticides. Plants were used against pests for their repulsive, contact or fumigating effects. Active molecules can vary from one family to another, within the same family and sensitivity can differ for one given insect from one stage to another. According to the studies reported by the authors [19], *Tephrosia vogelii* Hook f. has a repulsive and toxic action on *C. maculatus*, whereas *Blumea aurita* (L.) DC, which has no toxicity, has a strong repellent power over the same depreator. In addition, *Dracaena arborea* harvested in the Mono department in Benin is repulsive while the same species collected in the Borgou department is not effective.

Other researchers have taken a look at the chemical composition and biological properties of *Cymbopogon citratus* (DC.) Stapf (Poaceae), *Cymbopogon schoenanthus* (L.) Spreng. (Poaceae), *Cymbopogon giganteus* Chiov. (Poaceae), and *Eucalyptus citriodora* Hook. (*Myrtaceae*) in the fight against the depressor *Tribolium castaneum* [20] and *Callosobruchus maculatus* [19].

In the same perspective, other authors [21 - 23] assessed the toxicity of essential oils of *Dennettia tripetala*, *Uvariandron angustifolium*, *Premna angolensis*, *Premna quadrifolia*, *Polyalthia longifolia* and *Clausena anisata* against *Sitotroga cerealella*, a rice depreator. The chemical analysis by GC and GC / MS of these extracts showed that the *D. tripetala* essential oil was mainly composed of 2-Phenylnitroethane (52.6%), linalol (26.8%) and methyl eugenol (5.6%); that of *U. angustifolium* was dominated by geranial (44.9%), neral (32.1%) and geraniol (2.0%). The essential oil of *P. angolensis* contained 29 compounds representing 96.1% of oil against 42 compounds corresponding to 91% for the essential oil of *P. quadrifolia*. The majority compounds of these two oils (*P. angolensis* and *P. quadrifolia*) were β -caryophyllene (13.1%), (E) - β -

caryophyllene (13.5%), octen-3-ol (3.2% -28%), phytol (3.7% -4.9%), β -elemene (1.4% -21%), globulol (11.2%), germacrene-D (8.9%), α -humulin (2.9% -6.4%), α -pinene (5%), sabinene (3.7%), δ -cadinene (0 , 4% -3.3%) and linalool (3.3%). As for the essential oils of *Polyalthia longifolia* and *Clausena anisata*, the majority compounds were β -Caryophyllene (24.5%), Allo-Aromadendrene (13.5%), α -Zingiberene (9.4%), α -Humulene (8.5%) and α -Selinene (2, 6%) for *Polyalthia longifolia* oil and chavicol methyl oil (69.9%) for *Clausena anisata*. Oral toxicity assessment against *S. cerealella* at the laboratory by fumigation method in a glass jar closed at 29 ± 2 ° C and natural photoperiod with relative humidity of $70 \pm 10\%$ showed that these extracts have an insecticide and repulsive activity on the alucite. In addition, the toxicity of essential oils was also illustrated by the significant inhibition of insect emergence compared to the control groups without affecting the germination of treated rice grains.

These results suggest that studied volatile extracts can be used as a substitute for synthetic chemicals in paddy rice protection against *Sitotroga cerealella* and the reduction of food poisoning due to the use mainly of synthetic chemicals. Table 1 provides a summary of the chemical composition and the biological activity of a few species of plants.

Potential Food Applications of Vegetal Extractions in Conservation and Preservation of Public Sante

Antimicrobial properties are due to the essential oil fraction and secondary metabolites contained in the plants. The current trend of consumers to seek safe food and without food additives has led to a renewed interest in the Beninese scientists for these substances. In fact, for decades, studies have been conducted on the development of new applications and the exploitation of natural properties of plant extracts in the food sector.

Several extracts of plants have shown, in the laboratory, proven antimicrobial activity. But before they adoption as a food conservation agent, experimental results should be checked in the selected food. In general, the experimental results obtained in the model medium are confirmed on foods, but with slightly higher oils concentrations. Thus, the authors reported the efficacy of plant extracts for tomato [46], milk [47], milk waragashi cheese [48], peanut [49], fish [50], tchakpalo (local beer) [10], mango purse [51] and cowpea [52 - 53] conservation.

Table 1: Chemical and biological profiles of some plant species studied in Benin

Name of the plant / Family	Used parts	Identified chemical compounds	Sensitive microorganisms	MCB ou MIC (mgxmL-1)	Other activities detected	References
<i>X. Americana</i> (Olacaceae)	Trunk bark	phenolics, alkaloids sterols and terpenes	<i>Escherichia coli</i>	50 ¹	antiradical	[24]
<i>K.senegalensis</i>	Trunk	phenolics, alkaloids,	<i>Staphylococcus aureus</i>	0,78-6,25 ¹	antiradical	[25]
<i>P. kotschy</i> (Meliaceae)	bark	sterols and terpenes	<i>Escherichia coli</i> , <i>Salmonella typhi</i> et <i>Klebsiella pneumoniae</i>			
<i>C. quadrangularis</i> (Vitaceae)	Leafed stem	phenolics alkaloids, sterols and terpenes	<i>Salmonella typhi</i>	0,78 ¹		[8]
<i>D. oliveri</i> (Fabaceae)	Trunk bark	catechol	<i>Escherichia coli</i>	0,78 ¹	antiradical	[26]
<i>C. planchoni</i> (Cochlosperma-ceae)	Root bark	flavonoids, alkaloids	<i>Staphylococcus aureus</i> , <i>Escherichia coli</i> , <i>Salmonella typhi</i> et de <i>Klebsiella pneumoniae</i> ,	6,25-100 ¹	antiradical, insecticide	[27] [28]
<i>Eucalyptus citriodora</i> (Myrtaceae)	leaf	Isopulégol, citronellal et citronellol	<i>Staphylococcus aureus</i> , <i>Escherichia coli</i> , <i>Candida albicans</i>	0,63 à 1,25 ¹	antiradical	[29]
<i>S.longepedunculata</i> (Polygalaceae)	Root bark	salicylate méthyle	<i>Escherichia coli</i> , <i>Candida albicans</i>	0,4-12,79	antiradical	[7]
<i>D. africanum</i> (Apiaceae)		Methylstyrene, Mentatriene, Limonene	<i>Staphylococcus aureus</i> , <i>Escherichia coli</i> , <i>Candida albicans</i>	0,53-8,43		[30]
<i>C. odorata</i> (Asteraceae)	leaf	Germacrène, pregeijerène, geijerène, pinene	<i>Staphylococcus aureus</i> , <i>Escherichia coli</i>	1,28-5,11		[31]
<i>C. Rubens</i> (Asteraceae)	leaf	Myrcene, limonène, thujene	<i>Staphylococcus aureus</i> , <i>Escherichia coli</i> , <i>Candida albicans</i> ,	1,08-4,38		[32]
<i>H. suaveolens</i> (Lamiaceae)	Aerial part	phenolics, tannins,	<i>Plasmodium falciparum</i>		antiradical, antiplasmodial	[9]
<i>S. caudatum</i> (Poaceae)	stem	phenolics, tannins,				[33]
<i>O. canum</i> (Lamiaceae)	leaf	Terpinène ; linalol, terpinèn-4-ol, caryophyllène			Fungicide antimicrobial	[34]

Continued on next page

Table 1 continued

Name of the plant / Family	Used parts	Identified chemical compounds	Sensitive microorganisms	MCB ou MIC (mgxmL-1)	Other activities detected	References
<i>P. senegalensis</i> (<i>Polygonaceae</i>)	leaf	phenolics, alkaloids	<i>Staphylococcus aureus</i> , <i>Candida albicans</i> ,	5-20	antiradical	[35]
<i>P. butyracea</i> (<i>Clusiaceae</i>)	bark	Caryophyllène, humulène				[36]
<i>C. giganteus</i> (<i>Poaceae</i>)	leaf	cis-p-mentha-1(7), 8-dien-2-ol, trans-mentha-2,8-dien-1-ol limonène	<i>Escherichia coli</i> , <i>Staphylococcus aureus</i>	0,32-064	insecticide	[37]
<i>C. schoenanthus</i> (<i>Poaceae</i>)	leaf	piperitone, 2-carene	<i>Escherichia coli</i> , <i>staphylococcus aureus</i>	0,63	antiradical	[37]
<i>C. ambrosioides</i> (<i>Chenopodiaceae</i>)	leaf	terpinen, ascaridole, p-cymene, isoascaridole	<i>Escherichia coli</i> , <i>staphylococcus aureus</i>	1,56-6,69		[37]
<i>P. racemosa</i> (<i>Myrtaceae</i>)	leaf	eugenol, myrcene, chavicol, limonene, 1,8-cineole	<i>Escherichia coli</i> , <i>staphylococcus aureus</i> <i>Salmonella typhi</i> , <i>Candida albicans</i>	0,24-2, 24		[37]
<i>C. zeylanicum</i> (<i>Lauraceae</i>)	leaf	cinnamaldehyde, cinnamyl acetate, cinnamyl benzoate;	<i>Escherichia coli</i> , <i>staphylococcus aureus</i> , <i>Candida albicans</i>	0,2-1,6		[38]
<i>S. aethiopicus</i> (<i>Zingiberaceae</i>)	leaf					[39]
<i>A. difformis</i> (<i>Araceae</i>)	racine	ethyl acetate, Toluene, décadiyne, cymene, caryophyllène			antiradical	[40]
<i>P. biglobosa</i> (<i>Mimosaceae</i>)	leaf	ethyl acetate,			antiradical	[40]
<i>P. longifolia</i> (<i>Annonaceae</i>)	leaf	ethyl acetate,			antiradical	[40]
<i>N. laevis</i> (<i>Bignoniaceae</i>)	leaf	flavonoids, coumarins, tannins	<i>Salmonella typhi</i> , <i>Klebsiella pneumoniae</i> , <i>Escherichia coli</i>	0,32-1,25	antiradical	[41]
<i>P. Santalinoides</i> (<i>Fabaceae</i>)	leaf	flavonoids, tannins	<i>Salmonella typhi</i> , <i>Klebsiella pneumoniae</i> , <i>Escherichia coli</i>	0,32-1,25	antiradical	[41]

Continued on next page

Table 1 continued

Name of the plant / Family	Used parts	Identified chemical compounds	Sensitive microorganisms	MCB ou MIC (mgxmL-1)	Other activities detected	References
<i>Eucalyptus tereticornis</i> (Myrtaceae)	leaf	cymene, Crypton			insecticide	[42]
<i>Cochlospermum tinctorium</i> (Bixaceae)	leaf	cyclododecanone,			insecticide	[42]
<i>Citrus sinensis</i> (Rutacea)	leaf	sabinene, Limonene,	<i>Salmonella typhi</i> , <i>Klebsiella pneumoniae</i> , <i>Escherichia coli</i>		antibactérienne	[43]
<i>Melaleuca leucadendron</i>	leaf	Terpineol, cymene, nerolidol, ledol	cineole, pinene,		insecticide	[23]
<i>Aframomum sceptrum</i> (Zingiberaceae)	leaf	Pinene, Sabinene, myrcène, phellandrène, linalool, borneol, myrtenyl acetate, elemene, caryophyllene, humulene, germacrene				[22]
<i>Bridellia ferruginea</i> (Euphorbiaceae)	Bark of stem	flavonoids, alkaloids,	tannins,		antiradical	[44]
<i>Ceiba pentandra</i> (Bombacaceae)	root	flavonoids			antiradical	[44]
<i>Dennettia tripetala</i>	leaf	Phenylnitroethane, linalol, méthyleugénol			insecticides	[22]
<i>Uvariadendron angustifolium</i>	leaf	Géranial, le néral et le géranol			insecticides	[22]
<i>Pimenta racemosa</i> (Myrtaceae)	leaf	Mycenes, eugénol	chavicol,			[45]
<i>Hemizygia bracteosa</i> (Lamiaceae)	leaf	Known chemical profile				[13]

¹=calculated value for MICs; - Not determined

Thus, plant extracts, especially the essential oils extracted from acclimated aromatic plants in Benin, were first addressed from their chemical composition as well as their antibacterial and antifungal power against many pathogens and alteration strains of foods. In this context, the chemical composition and biological properties of essential oils extracted from the fresh leaves of *Lippia rugosa* A. Chev [11], *Cymbopogon citratus* [54 - 55] and *Ocimum scissimum* [47] have been investigated for their use in the conservation of food products. The results of these studies have shown that the essential oils extracted from *Lippia Rugosa* A. Chev, *Cymbopogon citratus* and *Ocimum gratissimum* collected in Benin, have interesting antimicrobial properties, offering new perspectives in the fight against microorganisms which contaminate foodstuffs. Better still, the essential oil extracted from the fresh leaves of *O. gratissimum* has properties of bio-conservation of the tomato, food of great importance for the Beninese population. The antiaflatoxinogenic power of these extracts was also invested in the protection of cereals and peanuts, in large post-harvesting systems against mold contamination and the production of mycotoxins [56 - 58]. The results show the antiaflatoxinogenic power of essential oils extracted from fresh leaves of *Lantana Camara*, *Congzoidate Ageratum*, *Ocimum Scissimum* and *Ocimum Canum* harvested in Benin against toxinogenic peanut millet.

In the same order, other investigations focused on the preservation of highly perishable food products such as fish products. Indeed, in Benin, fishing has a relatively significant place in the national socio-economic balance because it contributes approximately 3% to the Brut Interior Product (GDP). However, fish conservation is difficult because of the lack of appropriate adequate conservation and climatic and environmental conditions that contribute to its rapid degradation. To limit these losses, fumes are one of the conservation methods of the fish. However, despite the many efforts made for its conservation, smoked fish remains still very perishable due to increased microbial proliferation. In this context, the chemical composition of the essential oil of *Syzygium aromaticum* acclimated in Benin and its antimicrobial properties against smoke-free ingredients, especially those of the trailer (*Trachurus trachurus*) were invested on the one hand [59]. On the other hand, the comparative efficiency of essential oils for high antibacterial activity, extracted from aromatic plants, including *Ocimum gratissimum*, *Pimenta racemosa*, *Cymbopogon citratus* and *Mentha Piperita* was studied for their use as natural conservatives of very much perishable like fried fish [59], cheese [60] and fresh cow milk [48]. These plants have the advantage of being already part of the Beninese culinary traditions.

Still with a view to valorizing agro-resources and taking into account endogenous knowledge, the plant species *Hemizygia*

bracteosa (Benth) has been the subject of deep investigations. Indeed, in rural areas people use every day, local drinks from traditional production processes during festive celebrations. However, the technologies used as well as the production conditions still do not guarantee enough the health and organoleptic quality of these beverages. Thus, a diagnostic study has led to the status of endogenous knowledge of local beverages, as well as the use of plant extracts in the improvement their hygienic and organoleptic quality [10]. Similarly, the botanical species *Hemizygia bracteosa* (Benth), widely used in ethnomedicine in Benin, was studied for its biological properties in improving the sanitary and organoleptic quality of traditional *tchapalo* beer [61]. However, toxicity tests on the Wistar Albinic rat showed a toxicity at a dose of 3g / Kg of body weight on the liver [13].

These results provide new perspectives in the preservation of highly perishable foodstuffs in Benin and also illustrate the importance of using natural extracts of plants as an alternative in replacing commonly used synthetic chemicals.

Lastly, the biological properties of plant species *Spondias Mombin* L. and *Vitellaria paradoxa* Gaertn F. were also evaluated for use as a source of molecules to galactogenic properties [62]. They provide new perspectives in solving one of the many problems related to infant feeding.

However, there are some limits to the use of essential oils as food conservation agents, notably the flavoring power of some of them. However, dearomatization techniques exist and are more and more effective. Better, undesirable organoleptic effects may be limited by carefully selecting the essential oil according to the type of food concerned, but it is important to note that, in most cases, the oils concentrations used are so low, they do not alter the organoleptic qualities of food [63].

Table 2 shows effective thresholds (Minimum Inhibitor Concentration) of some essential oils for pathogens and an overview of studies showing antimicrobial activity of plant species in foods.

Conclusion and Perspectives

This work has made the inventory of research results in chemistry of biologically active substances in Benin. It appears that medicinal plants are a valuable heritage for Benin and still remain a significant source of medical care. Specific wealth and diversity of uses demonstrate the level of knowledge of medicinal plants and the dependence of the local population for primary health care. The research strives to identify the active ingredients in the perspective of developing improved traditional medicine. On the other hand, many natural properties of plant extracts make very promising conservation agents for the food

Table 2 Plant species, chemical compositions and potential application in agri-food of some plant species

Plant species	Chemical compositions (%)	Sensitive isolated strains (from food-stuffs)	Concentration of extracts tested ($\mu\text{L.mL}^{-1}$)	Probable Uses	References
<i>Cymbopogon citratus</i>	Myrcene (10,4), Neral (33), Geraniol (6,6), Geraniol (41,3),	<i>Saccharomyces cerevisiae</i> , <i>Aspergillus niger</i> , <i>Fusarium oxysporum</i> , <i>Penicillium camembertii</i>	500-2500	*Conservation of traditional <i>tchakpalo</i> beer * Preservation of the peuhl cheese <i>waragashi</i> * Preservation of cow's milk	[48][55-56]
<i>Ocimum gratissimum</i>	γ -terpinene (20,0), thymol (26,9), p-cymene (17,6), α -thujene (8,2).	<i>Aspergillus candidus</i> , <i>Penicillium camemberti</i> , <i>E. coli</i> , <i>S. aureus</i> , <i>Fusarium oxysporum</i> , <i>Fusarium graminearum</i> , <i>Fusarium poae</i> , <i>Aspergillus niger</i>	5-1600	* Preservation of tomato * Preservation of smoked fish	[47] [60]
<i>Pimenta racemosa</i>		<i>Aspergillus candidus</i> , <i>Penicillium camemberti</i>	5-10	Smoked fish	[60] [53]
<i>Hemizygia bracteosa</i>	(E)-b-Farnesene (67), trans-Nerolidol (6,2), b-Elémène (7.4)	<i>Penicillium citrinum</i> , <i>Aspergillus oryzae</i> , <i>Penicillium italicum</i> , <i>Penicillium citrinum</i> , <i>Aspergillus flavus</i> , <i>Aspergillus parasiticus</i>	0.5-1.5	Conservation of traditional <i>tchakpalo</i> beer	[62] [64]
<i>Mentha spicata</i>	carvone (66.57), limonène (16.33) and 1.8-cineole (7.22)	<i>Aspergillus parasiticus</i> , <i>Penicillium digitatum</i>	100-300*	Conservation of traditional beer <i>tchakpalo</i>	[65]
<i>Mentha piperita</i>	1,8-Cineole (6,5), <i>Menthone</i> (7,4), Menthol (46,7), neo-menthol (8,28), menthyl acetate (6,7),	<i>Aspergillus parasiticus</i> , <i>A. versicolor</i> and <i>Mucor spp</i>	0.50-2.5	Conservation of mango puree	[51]
<i>Tectona grandis</i>	Gallic tannins, catechic tannins, flavonoids, anthocyanins,	-	-	Food packaging (corn dough)	[66]

industry. Each plant extract has a variable biological activity depending on the microorganisms tested.

The use of extracts proves to be a relevant choice in the face of a specific risk of contamination or the need to reduce or replace synthetic chemical preservatives. In addition, their use in very small amounts could be envisaged due to their high efficacy. Their combined use with other conservation methods will definitely make the most important natural antimicrobial agents in the coming years to improve food life. Other properties of extracts, including antiparasitic, insecticide, antimicrobial and antiviral drugs are currently being studied to meet medical requirements, biological farming (developing biopesticides, producing animal supplements.).

In the medium and long term, this work could be an answer to the problem of antibiotics and their resistance, and to have a human and animal health application.

References

- [1]. Jiofack T, Fokunang C, Guedje N, Kemeuze V, Fongzossie E, Nkongmeneck B A, Tsabang N. Ethnobotanical uses of medicinal plants of two ethnoecological regions of Cameroon. *Int. J. Med. Med. Sci.* 2010; 2(3): 60-79.
- [2]. Laplante J. Le médicament aux frontières des savoirs humanitaires et autochtones. *Anthropologie et sociétés*, 2003; 27(2): 59-75.

- [3]. Salhi S, Fadli M, Zidane L, Douira A. Etudes floristique et ethnobotanique des plantes médicinales de la ville de Kénitra (Maroc). *Lazaroa*, 2010 ; 31 : 133.
- [4]. Lawin IF, Laleye OAF, Agbani O P. Vulnérabilité et stratégies endogènes de conservation des plantes utilisées dans le traitement du diabète dans les communes de Glazoué et Savè au Centre-Bénin. *Int. J. Biol. Chem. Sci.*, 2016 ; 10(3) : 1069-1085.
- [5]. Codjia JTC, Assogbadjo AE, Ekué MRM. Diversité et valorisation au niveau local des ressources végétales forestières alimentaires du Bénin. *Cahiers agricultures*, 2003 ; 12(5) : 321-331.
- [6]. Adomou AC, Yedomonhan H, Djossa B, Legba SI, Oumorou M, Akoegninou, A. Etude Ethnobotanique des plantes médicinales vendues dans le marché d'Abomey-Calavi au Bénin. *Int. J. Biol. Chem. Sci.* 2012; 6(2): 745-772
- [7]. Alitonou GA, Avlessi F, Tchobo FP, Noudogbessi JP, Tonouhewa A, Yehouenou B, Menut C, Sohounhloué KD. Chemical composition and biological activities of essential oils from leaves of *Cymbopogon giganteus* Chiov. and *Cymbopogon schoenanthus* (L.) spreng (Poaceae) from Benin. *Int. J. Biol. Chem. Sci.* 2012; 6 (4): 1819-1827.
- [8]. Koudoro YA, Agbangnan DCP, Yèhouénoù B, Tchobo FP, Alitonou GA, Avlessi F, Akoègninou A, Sohounhloué KCD. Chemical characterization and biological activities of extracts from two plants (*Cissus quadrangularis* and *Acacia polyacantha*) used in veterinary medicine in Benin. *J. Pharmacogn. Phytochem.* 2015; 3(6): 91-96.
- [9]. Medoatinsa SE, Agbangnan CP, Viwami F, Bogninou AGSR, Noudogbessi JP, Lagnika L, Ahissou H, Sohounhloué DCK. In vitro antiplasmodial and antioxidant activities of ethanolic and hydroethanolic extracts of *hyptis Suaveolens*, *World J Pharm Pharm Sci.* 2015; 4(08):101-112.
- [10]. Konfo CTR, Worou C N., Dahoué -Ahoussi E, Cakpochichi M, Oumaou MM, Sohounhloué DCK. Improvement of African traditional sorghum beers quality and potential application of its extracts for their stabilization : a review. *J. Microbiol. Biotechnol. Food Sci.* 2015 ; 5 (2): 190-196.
- [11]. Yehouenou B , Ahoussi E , Sessou , Alitonou GA, Toukourou F, Sohounhloué DCK. Chemical composition and antimicrobial activities of essential oils (EO) extracted from leaves of *Lippia rugosa* A. Chev against food pathogenic and adulterated microorganisms. *Afr. J. Microbiol. Res.* 2012; 6 (26): 5496-5505.
- [12]. Mamadou SR, Moussa I, Sessou P, Yehouenou B, Agbangnan P, Illagouma A, Abdoulaye A, Sohounhloué D, Ikhiriki K. Etude phytochimique, activités antiradicalaire, antibactérienne et antifongique d'extraits de *Sebastiania chamaelea* (L.) Müll.Arg. *J. Soc. Ouest Afr. Chim.* 2014 ; 037 :10-17.
- [13]. Chabi NW, Konfo TRC, Adjagba M, Moussedikou L, Ahoussi E, Dahouenon E, Layeye A, G aguidi C, Soumanou MM. Evaluation of the Toxicity of *Hemizygia ractea* (enth) lant U ed in Traditional Medicine for the Treatment of Diabetes Mellitus in Benin. *Am. J. Biomed. Res.* 2015: 3(3): 40-44.
- [14]. Houndje EM, Ogni CA, Noudeke N, Farougou S, Yousao AK, Kpodekon TM. Recettes ethno-vétérinaire à base de plantes médicinales utilisées pour le traitement de la fièvre aphteuse au Bénin. *International Journal of Biological and Chemical Sciences.* 2016;10(5):2090-107.
- [15]. Dassou HG, Ogni CA, Yédomonhan H, Adomou AC, Tossou M, Dougnon JT, Akoègninou A. Diversité, usages vétérinaires et vulnérabilité des plantes médicinales au Nord-Bénin. *International Journal of Biological and Chemical Sciences.* 2014;8(1):189-210.
- [16]. Ogni CA, Kpodékon MT, Dassou HG, Boko CK, Koutinhoun BG, Dougnon JT, Youssao AK, Yedomonhan H, Akoegninou A. Inventaire ethno-pharmacologique des plantes utilisées dans le traitement des pathologies parasitaires dans les élevages extensifs et semi-intensifs du Bénin. *International Journal of Biological and Chemical Sciences.* 2014;8(3):1089-102.
- [17]. Alain KY, Oronce DO, Boniface Y, Mhuro Y, Pascal AD, Paul TF, Alain AG, Akpovi A, Dominique SK. Free radical scavenging and antibacterial potential of two plants extracts (*Khaya senegalensis* and *Pseudocedrela kotschy*) used in veterinary pharmacopoeia in Benin.
- [18]. Fandohan P, Zoumenou D, Hounhouigan DJ, Marasas WF, Wingfield MJ, Hell K. Fate of aflatoxins and fumonisins during the processing of maize into food products in Benin. *International Journal of Food Microbiology.* 2005 Feb 15;98(3):249-59.
- [19]. Brice K, Adjou Euloge S, Edwige DA, Konfo TR, Christian AB, Dominique S. Problématique de la conservation du niébé (*Vigna unguiculata* (L), Walp) en Afrique de l'Ouest: étude d'impact et approche de solution. *Journal of Animal & Plant Sciences.* 2016;31(1):4831-42.
- [20]. Bossou DA, Ahoussi E, Ruysbergh E, Adams A, De Kimpe N, Avlessi F, Sohounhloué CKD, Mangelinckx S, Smaghe G. Characterization of volatile compounds from three *Cymbopogon* Species and *Eucalyptus citriodora* from Benin and their insecticidal activities against *Tribolium castaneum*. *Ind. Crops Prod.* 2015; 76: 306–317.
- [21]. Adjalien E, Sessou P, Bothon TDF, Dangou J, Odjo T, Figueredo G, Noudogbessi JP, Kossou D, Menut C, Sohounhloué D. Chemical composition and bioefficacy of *Dennet-*

tia tripetala and Uvariadendron angustifolium leaves essential oils against the angoumois grain moth, Sitotroga cerealella. Int. J. Biosci. 2014; 5(8): 161-172.

[22]. Adjalian E, Sessou P, Yehouenou B, Bothon TDF, Noudogbessi JP, Dansou K, Menut C, Sohounhloué D. Antioviposition and repellent activity of essential oil from melaleuca leucadendron leaf acclimated in bénin against the angoumois grain moth. Int. J. Biol. Pharm. Allied Sci. 2015; 4(2): 797-806.

[23]. Adjalian E, Sessou P, Alitonou GA, Noudogbessi JP, Kossou D, Menut C, Sohounhloué D. Ovicide and Larvicide Activities of the Essential Oils Extracted from Polyalthia longifolia and Clausena anisata on Sitotroga cerealella. Online J. Biol. Sci. 2017.

[24]. Koudoro YA, Yovo M, Yehouenou B, Agbangnan DCP, Tchobo FP, Alitonou GA, Sohounhloué CKD. Chemical study, antiradical and antibacterial potential of the extracts of Ximenia americana and Cussonia arborea of Benin. World J Pharm Sci. 2014; 2(12): 1626-1635.

[25]. Koudoro YA, Dedome O, Yehouenou B, Yovo M, Agbangnan DCP, Tchobo FP, Sohounhloué KCD. Free radical scavenging and antibacterial potential of two plants extracts (*Khaya senegalensis* and *Pseudocedrela kotschy*) used in veterinary pharmacopoeia in Benin. Elixir Int. J. 2014; 76: 28720-28726.

[26]. Koudoro YA, Wotto DV, Konfo TRC, Agbangnan DCP, Sohounhloué CKD. Phytochemical screening, antibacterial and anti-radical activities of Daniellia oliveri trunk bark extracts used in veterinary medicine against gastrointestinal diseases in Benin, Int. J. Adv. Res. 2015; 10 (3): 1190-1198.

[27]. Bossou AD, Mangelinckx S, Yedomonhan H, Boko PM, Akogbeto MC, De Kimpe N, Sohounhloué DC. Chemical composition and insecticidal activity of plant essential oils from Benin against Anopheles gambiae (Giles). Parasites & vectors, 2013; 6(1): 337.

[28]. Koudoro YA, Dedomè LSO, Yovo M, Agbangnan DCP, Tchobo FP, Alitonou GA, Avlessi F, Sohounhloué CKD. Caractérisation chimique, activités antiradicalaire et antibactérienne des extraits de l'écorce de racine de Cochlospermum planchonii du Bénin. IJIAS. 2014 ; 4 (7) : 1582-1594.

[29]. Koudoro YA, Dossa CPA, Yèhouénou BB, Tchobo FP, Alitonou GA, Avlessi F, Sohounhloué DC. Phytochemistry, antimicrobial and antiradical activities evaluation of essential oils, ethanolic and hydroethanolic extracts of the leaves of Eucalyptus citriodora hook from Benin. Scientific Study & Res. 2014; 15(1): 59.

[30]. Koudoro YA, Alitonou GA, Sossou DJ, Yehouenou B, Avlessi F, Menut C, Sohounhloué, D. Chemical Composition and biological Activities of Essential Oil from Benin

Diplolophium africanum Turcz stem leaves. J. Soc Ouest Afr. Chim. 2011; 032: 1-8.

[31]. Avlessi F, Alitonou GA, Djenontin TS, Tchobo F, Yèhouénou B, Menut C, Sohounhloué D. Chemical composition and Biological activities of the Essential oil extracted from the Fresh leaves of Chromolaena odorata (L. Robinson) growing in Benin. J. Biol. Sci. 2012; 1(3): 7-13.

[32]. Yehouenou B, Wotto V, Bankole H, Sessou P, Noudogbessi JP, Sohounhloué D. Chemical study and antimicrobial activities of volatile extracts from fresh leaves of cras-socephalum rubens (juss & jack) s. Moore against food-borne pathogens. Scientific Study & Res. 2010; 11(3): 341-349.

[33]. Agbangnan DCP, Tachon C, Bonin C, Chrostowka A, Fouquet E, Sohounhloué DC. Phytochemical study of a tinctorial plant of Benin traditional pharmacopoeia: the red sorghum (Sorghum caudatum) of Benin. Scientific Study & Res. 2012; 13(2): 121-135.

[34]. Adjou ES, Kouton S, Dahouenon-Ahoussi E, Sohounhloué DC, Soumanou MM. Antifungal activity of Ocimum canum essential oil against toxinogenic fungi isolated from peanut seeds in post-harvest in Benin. International Research Journal of Biological Sciences. 2012;1(7):20-6.

[35]. Bothon FTD, Gnanvossou D, Noudogbessi JP, Hanna R, Sohounhloué D. Bactrocera Cucurbitae response to four Cymbopogon species essential oils. J. Nat. Prod. 2013; (6): 147-155.

[36]. Alitonou G, Avlessi F, Sohounhloué CKD, Bessière JM, Menut C. Chemical and biological investigation on volatile constituents of Pentadesma butyracea Sabine (Clusiaceae) from Benin. J Essential Oil Res. 2010; 22: 138-140.

[37]. Alitonou GA, Avlessi F, Tchobo FP, Noudogbessi JP, Tonouhewa A, Yehouenou B, Menut C, Sohounhloué KD. Chemical composition and biological activities of essential oils from leaves of Cymbopogon giganteus Chiov and Cymbopogon schoenanthus (L.) spreng (Poaceae) from Benin. Int. J. Chem. Biol Sci. 2012; 6 (4): 1819-1827.

[38]. Yehouenou B, Sessou P, Houinsou RL, Noudogbessi JP, Alitonou GA, Toukourou F, Sohounhloué DCK. Chemical composition and Antimicrobial activities of Cinnamomum zeylanicum Blume dry Leaves essential oil against Food-borne Pathogens and Adulterated Microorganisms. Int. Res. J. Biol. Sci. 2012; 1(6): 18-25.

[39]. Noudogbessi JPA, Tchobo PF, Alitonou GA, Avlessi F, Soumanou M, Chalard P, Figueredo G, Chalchat JC, Sohounhloué DCK. Chemical Study of Extracts of Siphonochilus aethiopicus (Schweinf.) B.L. Burt (Zingiberaceae) from Benin. Asian J Chem. 2013; 25(15): 8489-8492.

- [40]. Akibou Ossen ML, Agbangnan DCP, Bossou A, Yedomonhan HP, Avlessi F. Activités antiradicalaires et étude des composés volatils de trois plantes de la médecine traditionnelle du Bénin: *Anchomanes difformis*, *Parkia biglobosa* et *Polyalthia*. IJIAS. 2014 ; 9(4) : 1609-1619.
- [41]. Bothon FT, Moustapha M, SophieBogninou G, Dossa CP, Yehouenou B, Medoatinsa SE, Noudogbessi JP, Avlessi F, Sohounhloue DC. Chemical Characterization and Biological Activities of *Newbouldia laevis* and *Pterocarpus Santalinoides* Leaves. Bull. Env. Pharmacol. Life Sci. 2014 Oct 11;3(11):09-15.
- [42]. Bossou AD, Mangelinckx S, Yedomonhan H, Boko PM, Akogbeto MC, De Kimpe N, Avlessi F, Sohounhloue DC. Chemical composition and insecticidal activity of plant essential oils from Benin against *Anopheles gambiae* (Giles). Parasites & vectors. 2013 Dec;6(1):337.
- [43]. Yovo M, Alitonou GA, Yedomonhan H, Tchobo F, Dedome O, Sessou P, Avlessi F, Menu C, Sohounhloué D. First report on chemical composition and antimicrobial activity of *Artabotrys velutinus* Scott-Elliot extracts against some clinical strains in Benin. Am. J. Appl. Chem. 2016;4:71-6.
- [44]. Bothon FT, Debiton E, Yedomonhan H, Avlessi F, Teulade JC, Sohounhloue DC. α -Glucosidase inhibition, antioxidant and cytotoxicity activities of semi-ethanolic extracts of *Bridellia ferruginea* benth. and *Ceiba pentandra* L. Gaerth from Benin. Research Journal of Chemical Sciences. 2012; 2(12): 31-36.
- [45]. Keke M, Yehouenou B, de Souza C, Sohounhloue D. Evaluation of hygienic and nutritional quality of peulh cheese treated by *Sorghum vulgare* (L) and *Pimenta racemosa* (Miller) extracts. Scientific Study & Research. 2009 Jan 1;10(1):29-46.
- [46]. Houinsou RD, Ahoussi E, Sessou P, Yèhouénu B, Sohounhloué D. Antimicrobial activities of essential oil extracted from leaves of *ocimum gratissimum* L. against pathogenic and adulterated microorganisms associated to tomato in Benin. International Journal of Biosciences (IJB). 2012;2(11):90-100.
- [47]. Degnon GR, Adjou ES, Metome G, Dahouenon-Ahoussi E. Efficacité des huiles essentielles de *Cymbopogon citratus* et de *Mentha piperita* dans la stabilisation du lait frais de vache au Sud du Bénin. International Journal of Biological and Chemical Sciences. 2016;10(4):1894-902.
- [48]. Sessou P, Farougou S, Kaneho S, Djenontin S, Alitonou GA, Azokpota P, Youssao I, Sohounhloué D. Bioefficacy of *Cymbopogon citratus* essential oil against foodborne pathogens in culture medium and in traditional cheese *wagashi* produced in Benin. International Research Journal Microbiology. 2012;3(12):406-15.
- [49]. Adjou ES, Aoumanou MM. Efficacité des extraits de plantes dans la lutte contre les moisissures toxigènes isolées de l'arachide en post-récolte au Bénin. Journal of Applied Biosciences. 2013;70(1):5555-66.
- [50]. Dègnon GR, Faton AN, Adjou ES, Noudogbessi JP, Dahouenon-Ahoussi E, Soumanou MM, Sohounhloue DC. Antifungal potential of Clove essential oil (*Syzygium aromaticum* L.) in the post-smoking preservation of mackerel (*Trachurus trachurus*) in Benin. International Research Journal of Biological Sciences. 2013;2(10):36-42.
- [51]. Adjou ES, René D, Edwige DA, Soumanou MM. Chemical composition and biological activity of essential oil from *Mentha piperita* L. leaves on the quality of mango puree during storage. Journal of Microbiology, Biotechnology & Food Sciences. 2017 Oct 1;7(2):97-100.
- [52]. Rose de Lima FH, Adjou ES, Ahoussi ED, Sohounhloué DC, Soumanou MM. Caractéristiques biochimique et sensorielle du niébé (*Vigna unguiculata*) conservé au moyen des huiles essentielles extraites de plantes de la famille des Myrtaceae [Biochemical and sensorial Characteristics of cowpea (*Vigna unguiculata*) stored with essentials oils extracted from plants of Myrtaceae family]. International Journal of Innovation and Applied Studies. 2014 Nov 1;9(1):428.
- [53]. Kpatinvoh B, Adjou ES, Dahouenon-Ahoussi E, Konfo TR. Ch., Atrevey BC, Sohounhloue D. Problématique de la conservation du niébé (*Vigna unguiculata* (L), Walp) en Afrique de l'Ouest: étude d'impact et approche de solution. Journal of Animal & Plant Sciences. 2016;31(1):4831-42.
- [54]. Dahouenon-Ahoussi E; Dègnon GR, Adjou ES, Atodjinou V, Sohounhloue CKD. Amélioration de la technologie de fabrication de la bière à base de matières amylicées locales : utilisation combinée du sorgho (*Sorghum bicolor*) et de la banane (*Musa acuminata*). REMISE. 2012 ; 6 (1): 67-83.
- [55]. Konfo C, Dahouenon-Ahoussi E, Sessou P, Yehouenou B, Djenontin S, De Souza C, Sohounhloué D. Stabilization of local drink "Tchakpalo" produced in Benin by addition of essential oil extracted from fresh leaves of *Cymbopogon citratus*. Int Res. J. Biol. Sci. 2012; 1(8): 40-49.
- [56]. Adjou ES, Dahouénon-Ahoussi E, Degnon R, Soumanou MM, Sohounhloue CKD. Bioefficacy of essential oil of *Lantana camara* from Benin against the growth of fungi and aflatoxin production. JRAA. 2012; 1(4): 112-12.
- [57]. Adjou ES, Kouton S, Dahouenon-Ahoussi E, Soumanou MM, Sohounhloue CKD. Effect of essential oil from fresh leaves of *Ocimum gratissimum* L. on mycoflora during Storage of Peanut in Benin, Mycotoxin Res. 2013; 29(1): 29-38.

[58]. Adjou ES, Dahouenon-Ahoussi E, Degnon R, Soumanou MM, Sohounhloue CKD. Investigations on bioactivity of essential oil of *Ageratum conyzoides* L., from Benin against the growth of fungi and aflatoxin production. *Int. J. Pharm. Sci. Rev. Res.* 2012; 13(1): 143-148.

[59]. Dègnon RG, Faton AN, Adjou ES, Tchobo FP, Ahoussi DE, Soumanou MM, Sohounhloue DCK. Efficacité comparée des huiles essentielles de deux plantes aromatiques dans la conservation post-fumage du Chinchard (*Trachurus trachurus*). *J. Anim. Plant Sci.* 2013 ; 19(1): 2831-2839.

[60]. Zannou A, Konfo TRC, Gbaguidi ANM, Ahoussi-Dahouenon E. Antimicrobial activity of extracts from *Cymbopogon citratus* L. and of *Mentha spicata* L. against fungal and bacterial strains isolated from peuhl's cheese (Waragashi) produced in Benin. *Int J. Adv Res.* 2015; 3(10): 1684-1695.

[61]. Konfo CT, Chabi NW, Agbadjizo J, Dahouenon-Ahoussi E, Soumanou MM, Sohounhloue DC. Influence de la feuille de *Hemizygia bracteosa* (Benth) sur la qualité de la bière du sorgho " tchakpalo" produite au Bénin. *International Journal of Innovation and Applied Studies.* 2014 Aug;7(2):453-63.

[62]. Guénolé AC, Delphin KO, Edwige AH, Sylvie AH. Effects of Leaves Extract from *Spondias Mombin* L. and *Vitellaria Paradoxa* Gaertn F. on West African Dwarf (WAD) Sheep Performance in Republic of Benin. *Journal of Animal Science Advances.* 2013;3(2):74-82.

[63]. Oussalah M, Caillet S, Saucier L, Lacroix M. Inhibitory effects of selected plant essential oils on four pathogen bacteria growth: *E. coli* O157:H7, *Salmonella typhimurium*, *Staphylococcus aureus* and *Listeria monocytogenes*. *Food Control.* 2007; 18 (5): 414-420.

[64]. Kpoviessi S, Agbani P, Gbaguidi F, Gbénuou J, Sinsin BA, Accrombessi G, Quetin-Leclercq J. Seasonal variations of volatile constituents of *Hemizygia bracteosa* (Benth.) Briq. aerial parts from Benin. *Comptes Rendus Chim.* 2016; 19(7): 890-894.

[65]. Dahouenon-Ahoussi E, Adjou ES, Dègnon G, Konfo RTC, Dahoué R, Soumanou MM, Sohounhloue CKD. Efficiency of the essential oil of *Mentha spicata* L. in the stabilization of sorghum beer "tchakpalo" produced in Benin. *Elixir Int. J.* 2017; 104: 46028-46033.

[66]. Zannou A, Gbaguidi MA, Ahoussi-Dahouenon E. Chemical characterization of *Tectona grandis*, *Thalia geniculata*, *Lasiomorpha senegalensis* and of *Musa acuminata* used as food packing's in Benin. *Journal of Pharmacognosy and Phytochemistry.* 2016 Sep 1;5(5):79.