



Evaluation of hepatoprotective potential of leaf and leaf callus extracts of *Anisochilus carnosus* (L) wall.

Nissar Ahmad Reshi^{1*}, Sudarshana Mysore Shankarasingh² and Girish Vasanaika Hodiya²

Abstract

The study was carried out to evaluate the hepatoprotective activity of leaf and leaf callus extracts of *Anisochilus carnosus* (L) Wall. against alcohol induced toxicity using HepG2 cell line. Leaf explants were cultured on Murashige and Skoog solid medium supplemented with different growth regulators. Prior to the determination of hepatoprotective property leaf and leaf callus extracts were subjected to the toxic dose study. The degree of hepatoprotection of extracts was determined by measuring cell viability percentage by MTT assay. The preliminary phytochemical analysis of leaf and leaf callus extracts was carried out by qualitative analysis. Maximum percentage of callus formation (98%) was obtained in MS medium fortified with 3 mg/l 2,4-D. HepG2 cells were pretreated with the different concentrations (below toxic dose) of leaf and leaf callus extracts for 72 hours followed by alcohol intoxication. Results revealed that ethanolic leaf extract pretreated HepG2 cells show 94% cell viability compared to the standard silymarin pretreated HepG2 cells which showed 81% cell viability. Leaf callus extracts also exhibited significant hepatoprotective activity where ethanolic callus extract pretreated HepG2 cells showed 86% viability after intoxication with alcohol. Results revealed that HepG2 cell viability percentage is dose dependent. Phytochemical studies revealed the presence of different secondary metabolites in leaf and leaf callus extracts. The bio-efficacy study confirms the presence of secondary metabolites of hepatoprotective nature in leaf and leaf callus of *A. carnosus*.

Introduction

From last few decades medicinal plants have become the focus of intense study for the validation of their traditional uses. In India, there are almost 8000 medicinal plant species which comprise approximately 50% of the higher flowering plants [1]. Many efforts have been made to unravel the new hepatoprotective drugs of plant origin. Pharmaceutical companies spend considerable time and money in developing therapeutics based on natural products. It has been reported that about 170 phytoconstituents isolated from 110 plants belonging to 55 families possess hepatoprotective activity [2]. The rising incidence of Alcohol Liver Diseases (ALD), one of the most serious consequences of the chronic alcohol abuse and is the second leading cause of death among all liver diseases [3], has necessitated a search for

the new hepatoprotective drugs. Alcohol induces a number of adverse metabolic changes in liver. Excessive consumption of alcohol for a long time leads to the steatosis, alcoholic hepatitis and cirrhosis which results in weight and volume change of liver [4]. Studies on human HepG2 cell lines have shown in earlier studies that ethanol is cytotoxic and is apoptotic in nature predominantly in liver [5]. There are no satisfactory remedies available for liver diseases; hence search for effective hepatoprotective drugs from natural products is continued [6]. Plants have been source of diverse medicinal therapeutic agents. Only a small number of traditionally used medicinal plants have been scientifically evaluated for the hepatoprotective property [7]. Screening of callus extracts for biological activities paves a way for the mass production of biologically viable secondary metabolites and thus provides an alternative for the *in vivo* plant material and thereby minimising any possible threat to the natural habitat of medicinal plants. *A. carnosus* (L) Wall. is an annual herb, 30-60 cm tall, robust and branched with fleshy leaves. It grows in the regions of

*Correspondence: nissarreshi@gmail.com

¹Department of Botany, School of Science, Sandip University, Nashik, 422 231, Maharashtra, India

Full list of author information is available at the end of the article.

Received: 27 Jun 2017, Accepted: 17 Oct 2017



Hong Kong, Macao, Shanghai, Tianjin, India, Myanmar and Sri Lanka. The plant has been traditionally used for the treatment of gastrointestinal disorders, respiratory disorders, cough, cold and fever [8]. Its popular herbal preparation together with *Ocimum basilicum*, *Mentha piperita* and *Alpinia galanga* is used against the symptoms of influenza, dermatitis and the slight illness that derives from the bites of bugs [9]. Essential oils have been extracted by hydrodistillation from the leaves and have been reported to be antimicrobial in nature [10]. The plant has been reported to have anti-inflammatory activity [11], antiulcer activity [12], antifungal property [13] and anticancer property [14]. In the present investigation an attempt was made to screen leaf and leaf callus extracts of *A. carnosus* for hepatoprotective activity against ethanol induced toxicity using HepG2 cell lines.

Materials and Methods

Collection of plant material and callus induction

A. carnosus plants were collected from the Nilgiri Hills, Western Ghats of Tamil Nadu, India and are being maintained in the medicinal plant garden of Department of Studies in Botany, University of Mysore, Mysuru. Healthy leaf explants were collected from the mother plant and were washed under running tap water for ten minutes to remove the soil and dust particles followed by treatment with fungicide bavistin (1% w/v) for 5 minutes. Explants were then washed thrice with sterile double distilled water and were treated with mercuric chloride (0.01% w/v) for 5 minutes followed by washing thrice with sterile double distilled water. Inside laminar airflow, leaf explants were cut into small pieces (1 sq.cm) and were inoculated onto the MS medium containing 3% sucrose, 0.08% agar and fortified with different concentrations of auxins. Prior to autoclaving at 121°C for 15 min, pH of the medium was adjusted to 5.8 using 1N NaOH and 1N HCl. The culture flasks were maintained in the incubation chamber under a 16 hours photoperiod at light intensity of $25\mu\text{mol/s}^2/\text{m}^2$ for 4 weeks. Each experiment was performed with five replicate and repeated thrice. The callus cultures were maintained for the period of five months and were periodically subcultured with 2-3 weeks of interval onto the fresh MS medium for callus proliferation. Subsequently, the callus was harvested at the transfer age of 3 weeks, kept in hot oven at 60 °C for 24 hours till a constant dry weight was obtained and the callus was then further exploited for extraction and hepatoprotective evaluation. Dry leaves and the *in vitro* leaf derived callus was coarse powdered using electric homogenizer. Aqueous extraction was carried out by mixing leaf and leaf callus coarse powder separately with deionised water in the ratio of 1:5 (w/v) in conical flask and allowed to settle in an oven at 50°C for 72 hours with occasional shaking. After 72 hours the ex-

tracts were filtered using Whatman filter paper [15]. The filtrate was lyophilised to dryness and stored in vials at 5°C for further use. Solvent extraction was carried out by taking 25 grams of dry leaf and callus coarse powder and filled in a thimble separately and extracted sequentially with 200 ml of petroleum ether, chloroform, ethyl acetate, ethanol and methanol in Soxhlet extractor for 48 hours. The solvent extracts were concentrated under reduced pressure and were stored at 5°C in vials for further use. All the leaf and leaf callus extracts were dissolved in 1% DMSO for the bio-efficacy evaluation [16]. HepG2 (Normal Human Liver Cell Line) cell culture was procured from National Centre for Cell Sciences (NCCS), Pune, India. Stock cells of HepG2 cell line were subcultured as monolayers in DMEM (Dulbecco's Modified Eagle's Medium) supplemented with 10% FBS (Fetal Bovine Serum) and penicillin (100 µg/ml) [17]. Prior to the screening of hepatoprotective activity, toxic dose studies of extracts (both leaf and leaf callus) and silymarin were carried out by MTT assay (3-(4,5-dimethylthiazole-2 yl)-2,5 diphenyl tetrazolium bromide). The IC₅₀ value of ethanol was also calculated by MTT assay [2]. HepG2 cells were subcultured, after 72 hours medium supernatant was flicked off, cell monolayer was trypsinized and the cell count was adjusted to the 1.0×10^5 cells/ml. To each well of the 96 well plate 200 µl of diluted cell suspension (1×10^5 cells approximately) was added and the plates were incubated for 72 hours at 37°C and 5% CO₂ for 24 hours. After 72 hours when monolayer of cells was formed, medium supernatant was flicked off and the 200 µl of DMEM containing leaf and leaf callus extracts in concentrations below toxic dose was added to each well containing cell monolayer. The plates were then incubated for 72 hours at 37°C and 5% CO₂. This boosting of cells with the different extracts is also known as pre-treatment protocol. After 72 hours, the medium containing extracts was flicked off and DMEM containing ethanol at 150 mM concentration was added to the each well containing extract and silymarin pretreated cell monolayers. The plates were again incubated for 72 hours at 37°C and 5% CO₂. After 72 hours of incubation, medium containing ethanol was flicked off and 100 µl of 5% MTT reagent in DMEM was added to each well and the plates were again incubated at 37°C for 3 hours. After 3 hours, the supernatant was discarded and 100 µl of solubilisation solution (1% DMSO) was added to each well and the plates were gently shaken in gyratory shaker to solubilise the formed formazan [18]. The absorbance was then measured by using microplate reader at 630 nm and the percentage of growth inhibition was calculated using formula given as under:

$$\text{Growth inhibition(\%)} = \frac{\text{Mean OD of the test group}}{\text{Mean OD of control group}} \times 100$$

All the assays were carried out in triplicates and the data analysis was carried out by using SPSS, DMRT. Preliminary phytochemical analysis of leaf and leaf callus extracts was carried out by the method described by Sinha et al [19].

Results

Callus induction was observed along the cut edges of leaf segments after two weeks of inoculation on MS medium supplemented with auxins. Callus induction percentage varied with respect to the concentration of the growth regulator used and the age of the explant. Young explants cultured on MS medium augmented with 2,4-D (1 mg/l – 3 mg/l) showed high percentage of callus induction and proliferation. IBA and NAA supplemented medium induced hard and compact callus with low percentage of proliferation (**Table 1**). Cytotoxic study of leaf and leaf derived callus extracts of *A. carnosus* was carried out to standardise extract concentrations and evaluate their hepatoprotective activity. Results revealed that *A. carnosus* (both leaf and leaf callus) above 225 µg/ml concentration is toxic to the HepG2 cells where the cell viability percentage was reduced to 40%. MTT assay for cytotoxicity of standard drug silymarin revealed that concentration above 75 µg/ml is toxic to the cells and hence 75 µg/ml was used as the test concentration for the subsequent study. IC₅₀ value of ethanol was reported to be 150 mM (0.69% ethanol). Cytotoxic effect of different concentrations of leaf, leaf callus and silymarin is presented in **Fig.1**.




Figure 1 Cytotoxic effect of silymarin, leaf extracts and leaf callus extracts of *A. carnosus* on HepG2 cells.

Change in cell viability percentage on ethanol induced toxicity in HepG2 cells pretreated with different leaf and leaf callus extracts was carried out by MTT assay. Ethanol (150 mM) intoxicated HepG2 cells showed a percentage viability of 50% after 72 hrs. *A. carnosus* leaf and leaf callus extracts were tested at

100 µg/ml and 200 µg/ml concentration for their hepatoprotective activity. Out of the different extracts of leaf and leaf callus, polar solvent extracts like, aqueous, ethanol and methanol pretreated cells showed significant viability percentage after intoxication with ethanol. HepG2 cells pretreated with leaf and leaf callus extracts showed a dose dependent increase in percentage viability. The cell viability ranged from 60% to 94% in leaf extracts and 58% to 86% in leaf callus extract pretreated cells.

The maximum cell viability percentage (94%) was reported in ethanolic leaf extract pretreated HepG2 cells at 200 µg/ml. Out of the callus extracts, 86% cell viability was reported in ethanolic callus extract pretreated HepG2 cells. Silymarin pretreated HepG2 cells showed maximum 81% cell viability. The cell viability percentage of HepG2 cells pretreated with leaf and leaf callus extracts is presented in **Table 2 & 3**.

Preliminary phytochemical analysis of leaf and leaf callus extracts revealed the presence of Alkaloids, flavonoids, saponins, phytosterols, phenols, terpenoids, triterpenoids and sterols.

Discussion

The study was aimed to evaluate the hepatoprotective property of leaf and leaf callus extracts of *A. carnosus*. Effect of auxins on callus induction has been reported earlier by many authors [20–22]. In our study, 2,4-D supplemented MS medium was found to induce maximum callusing in leaf segments. Our results are directly in concurrence with the earlier findings reported by Rout et al (1999) [20] and Manickam et al., (2000) in *Plumbago zeylanica* and *Withania somnifera* [23] respectively. Drug, whether synthetic or natural, if consumed in excess dose damages the liver cells. This is the reason; toxic dose study of drug is carried out. In the present study, cytotoxic study of *A. carnosus* leaf and leaf callus extracts was carried out by MTT assay before the extracts were subjected to the hepatoprotective activity. The principle involved is the cleavage of tetrazolium salt into its blue coloured derivative formazan; the process takes place in the inner mitochondrial membrane where an enzyme succinate dehydrogenase cleaves the tetrazolium salt. Only the living cell mitochondria reduces the MTT to coloured formazan, therefore the concentration of the dye is directly proportional to the number of metabolically active cells. The MTT assay revealed that concentration of *A. carnosus* leaf and leaf callus above 225 µg/ml kill 60% of the cells. Toxic dose studies of the plant extract lays the foundation for the formulation of any type of herbal remedy [24]. In the present investigation, ethanol was used as hepatotoxicant. Ethanol induced live injury is encompassed by wide spectrum of lesions, the most characteristic being alcoholic steatosis (fatty liver), alcoholic hepatitis, alcoholic fibrosis and cirrhosis [25]. Results revealed that ethanol treated

Table 1 Influence of auxins on callus induction from the leaf segments of *A. carnosus*

MS medium+Plant Growth Regulator mg/l	Mean percentage of callus induction	MS medium+Plant Growth Regulator mg/l	Mean percentage of callus induction
2,4-D		IAA	
0.5	68	0.5	43
1.0	86	1.0	47
1.5	90	1.5	56
2.0	90	2.0	62
2.5	92	2.5	64
3.0	98	3.0	70
3.5	72	3.5	76
4.0	70	4.0	79
4.5	74	4.5	82
5.0	68	5.0	78
IBA		NAA	
0.5	32	0.5	30
1.0	42	1.0	36
1.5	58	1.5	42
2.0	58	2.0	46
2.5	62	2.5	53
3.0	68	3.0	58
3.5	72	3.5	60
4.0	65	4.0	63
4.5	60	4.5	68
5.0	62	5.0	68

*All treatments with 6 replicates and were repeated thrice. Callusing percentage: (Number of explants callused/Total number of explants inoculated) x 100.

Table 2 The cell viability percentage of the HepG2 cells pretreated with leaf extracts of *A. carnosus*

Leaf Extracts	Concentration ($\mu\text{g/ml}$)	% cell viability
Aqueous	100	82 \pm 0.32
	200	85 \pm 0.26
Ethanol	100	88 \pm 0.51
	200	94 \pm 0.56
Methanol	100	78 \pm 0.57
	200	86 \pm 0.38
Ethyl acetate	100	73 \pm 0.20
	200	76 \pm 0.35
Petroleum ether	100	60 \pm 0.31
	200	60 \pm 0.32
Chloroform	100	60 \pm 0.38
	200	60 \pm 0.42
Silymarin	75	81 \pm 0.53
Control	Only cells	100

*All the MTT assays were repeated thrice. Each value represents Mean \pm S.D. Statistical analysis done by SPSS

Table 3 The cell viability percentage of the HepG2 cells pretreated with leaf callus extracts of *A. carnosus*

Callus Extracts	Concentration ($\mu\text{g/ml}$)	% cell viability
Aqueous	100	70 \pm 0.67
	200	74 \pm 0.31
Ethanol	100	78 \pm 0.20
	200	86 \pm 0.43
Methanol	100	80 \pm 0.43
	200	83 \pm 0.21
Ethyl acetate	100	62 \pm 0.42
	200	66 \pm 0.32
Petroleum ether	100	63 \pm 0.45
	200	65 \pm 0.40
Chloroform	100	58 \pm 0.23
	200	60 \pm 0.45
Silymarin	75	81 \pm 0.53
Control	Only cells	100

*All the MTT assays were repeated thrice. Each value represents Mean \pm S.D. Statistical analysis done by SPSS.

groups of HepG2 cells showed a drastic decrease in the cell viability when compared to the HepG2 cell pretreated with the leaf and leaf callus extracts. Increase in the percentage of cell viability in pretreated HepG2 cells indicates that the cells get boosted up upon treatment with extract and does not allow oxidation to take place upon intoxication with ethanol. In our study, HepG2 cells pretreated with leaf and leaf callus extracts showed percentage viability more than that of positive control silymarin. The cell viability relies on the structure of membrane and any dam-

age to cell membrane causes leakage of the cellular enzymes and consequently a cell death [2]. Results revealed that HepG2 cells pretreated with aqueous, ethanol and methanol leaf and leaf callus extracts of *A. carnosus* show a dose dependent increase in the cell viability. The dose dependent cytoprotection has been reported in other plants like *Cassia roxburghii* [26], *Polygonum multiflorum* [27], *Andrographis paniculata* [28] and *Rumex vesicarius* [29]. Kanchana and Jayapriya, (2013) [30] described in their study that the increase in the cell viability percentage may

be a consequence of membrane stabilisation boosted by phytochemicals and they further demonstrated that the plant extracts elevate the tissue antioxidant defence enzymes and thus tackles oxidative stress. Several researchers have concluded that the antioxidant activity of plant extract is generally attributed to the presence of phenolic compounds [31, 32]. Preliminary qualitative phytochemical studies revealed the rich presence of phenolic compounds in leaf and leaf callus extracts which could possibly be attributed to hepatoprotective nature of the extracts. Callus cultures have proved to be of great importance in alternative medicine in recent times as it paves a way for the establishment of callus cell lines for the mass production of the pharmacologically and biologically active secondary metabolites. Our results revealed that leaf callus extracts of *A. carnosus* possess hepatoprotective activity against ethanol induced toxicity in HepG2 cells. The chemical constituents of leaf and leaf callus extracts may have interrupted the reaction of ROS with cell proteins and nucleic acids and thus preventing the formation of adducts by acting as scavengers and thereby stabilising the cell membrane resulting in increased cell viability. Our inference is in concurrence with the earlier studies carried out by Santhosh et al. [33], Kamel et al [34] and Bhagyashree et al [35].

Conclusion

Ethanol induces production of reactive oxygen species (ROS), leading to huge oxidative stress which damages the liver cells. *A. carnosus* leaf and leaf callus extracts have shown the dose dependant hepatoprotection against ethanol. The results suggest the presence of active phytoconstituents in leaf and leaf callus extracts which strengthens antioxidant defence in cells and thus minimising the chances of production of free radicals. Our study further confirms the hepatoprotective property of leaf callus, which adds the extra feather to our study as the callus cultures could be used for the mass production of desired secondary metabolites and may also be used as replacement for the *in vivo* plant material and thus minimising the chances of threat to natural population of medicinal plants. Furthermore phytochemical analysis needs to be carried out to isolate and characterise the bioactive compounds from leaf and leaf callus extracts with hepatoprotective activity and for further authentication pharmacological studies also need to be carried out.

Conflict of interest

Authors declare that they do not have any conflict of interest.

Acknowledgments

Authors are grateful to the University of Mysore for providing all the necessary facilities to carry out this research work.

Author details

¹Department of Botany, School of Science,, Sandip University, Nashik, 422 231, Maharashtra, India. ²Department of Studies in Botany, University of Mysore, Mysuru, Karnataka, India.

References

- [1] Johnson M, Wesely E, Kavitha M, Uma V. Antibacterial activity of leaves and inter-nodal callus extracts of *Mentha arvensis* L. *Asian Pacific journal of tropical medicine*. 2011;4(3):196–200.
- [2] Ramoji A, Kuppast I, KLMankani. Protective effect of *Paracalyx scariosa* against ethanol induced hepatic toxicity in BRL 3A cell lines. *International Research Journal of Pharmacy*. 2014;5(6):495–501.
- [3] Neuman M. Cytokines - central factors in alcoholic liver disease. *Alcohol Research Health*. 2003;27(10):307–316.
- [4] Kumar V, Cotran R, Robbins S. Robbins basic pathology, 7th Ed. Philadelphia: W.B. Saunders; 2003.
- [5] Wu D, AICaderbaum. Ethanol induced apoptosis to stable HepG2 cell line expressing human cytochrome P-4502E1. *Alcoholism: Clinical and Experimental Research*. 1993;23(9):67–76.
- [6] Panda N, Banerjee S, Mandal N, Sahu N. Pragnane glycosides. *Natural Product Communications*. 2006;7(1):665–669.
- [7] GE T, WC E. Pharmacognosy. vol. 1983. London: Balliere Tindall Press; 1983.
- [8] Kamble S, More T, Patil S, Pawar S, Bindurani R, Bodhankar S. Plants used by the tribes of Northwest Maharashtra for the treatment of gastrointestinal disorders. *Indian Journal of Traditional Knowledge*. 2008;7(2):321–325.
- [9] Subramanian S, Nair A. Flavonoids of the leaves of *Mentha spicata* and *Anisochilus carnosus*. *Phytochemistry*. 1972;11(1):452–453.
- [10] Senatore F, Francesca L, Venza F, Bruno M, Napolitano F. Composition and antibacterial activity of the essential oil of *Anisochilus carnosus*- a Tamil plant acclimatized in Sicily. *Flavour and Fragrance Journal*. 2003;18(8):202–204.
- [11] Grover J, Adiga G, Vata V, Rathi S. Extracts of *Anisochilus carnosus* prevent development of experimental inflammation. *Journal of Ethanopharmacy*. 2001;7(8):159–164.
- [12] Mohammed A, Kumar R, Santosh HY, Nagashruthi M. Antiulcer activity of *Anisochilus carnosus* leaf extracts in pylorus ligation rats. *Indian Drugs*. 2008;45(12):979–983.

- [13] Kulandhaivel M, Sooraj S, Nathiya K, Palaniswamy M. Antifungal activity and phytochemical screening of *Anisochilus carnosus* and *Melaleuca alternifolia* against opportunistic pathogen *Candida albicans*. *International Journal of Pharmaceutical and Biological Archives*. 2011;2(6):1668–1670.
- [14] Muthuraman M, Santharam L, Ariraman S, Pemaiah B. Studies on anticancer and antimicrobial efficacy of *Anisochilus carnosus* extracts. *International Journal of Pharmacy and Pharmaceutical Sciences*. 2012;4(2):132–135.
- [15] JB H. *Phytochemical Methods-A guide to modern techniques of plant analysis*. Chapman and Hall, London; 1998. p. 182–190.
- [16] Escara J, Hutton R. Thermal stability and renaturation of DNA in dimethyl sulfoxide solutions: acceleration of the renaturation rate. *Biopolymers*. 1980;19(7):1315–1327.
- [17] Thabrew M, Hughes R, Mcfarlane I. Screening of hepatoprotective plant components using a HepG2 cell cytotoxicity assay. *Journal of Pharmacy and Pharmacology*. 1997;49(11):1132–1135.
- [18] H O, HS L, T K, KY C. Furocoumarins from *Angelica dahurica* with hepatoprotective activity on tacrine-induced cytotoxicity in Hep G2 cells. *Planta Medica*. 2002;68(5):463–464.
- [19] Sharma A, Gangwar M, Kumar D, Nath G, Sinha A, Tripathi Y. Phytochemical characterization, antimicrobial activity and reducing potential of seed oil, latex, machine oil and presscake of *Jatropha curcas*. *Avicenna Journal of Phytomedicine*. 2016;6(4):366–375.
- [20] Rout GR, Saxena C, Samantaray C, Das P. Rapid plant regeneration from callus cultures of *Plumbago zeylanica*. *Plant Cell Tissue and Organ Culture*. 1999;56:47–51.
- [21] Jain S, Jian R, Vlietinck J. In vitro and in vivo antimicrobial efficacy of *Mimosa hamate*. *Indian Journal of Biotechnology*. 2004;3:271–273.
- [22] Shariff N, Sudarshana M, Umesha S, Hariprasad P. Antimicrobial activity of *Rauvolfia tetraphylla* and *Physalis minima* leaf and leaf callus extracts. *African Journal of Biotechnology*. 2006;5:946–950.
- [23] Manickam V, Elangomadhavan R, Antonisamy R. Regeneration of Indian ginseng plantlets from stem callus. *Plant Cell Tissue and Organ Culture*. 2000;14:55–58.
- [24] Ibrahim M, Khaja N, Aara A, Khan A, Habeeb A, Devi P, et al. Hepatoprotective activity of *Sapindus mukorossi* and *Rheum emodi* extracts. *World Journal of Gastroenterology*. 2008;14(6):2566–2571.
- [25] OL K. Alcoholic liver disease: Pathobiological aspects. *Journal of Hepatology*. 1995;23(4):7–15.
- [26] Arulkumaran K, Rajasekaran A, Ramasamy R, Jegadeesan M, Kavimani S, Somasundaram S. *Cassia roxburghii* seeds protect liver against toxic effects of ethanol and carbontetrachloride in rats. *International Journal of Pharma Tech Research*. 2009;1(2):273–246.
- [27] HB L, YY H, CS W. Hepatoprotective activity of fresh *Polygonum multiflorum* against HepG2 hepatocarcinoma cell proliferation. *Journal of Food and Drug Analysis*. 2011;19(1):246–273.
- [28] Sivaraj A, Vinothkumar P, Sathiyaraj K, Sundaresan S, Devi K, Senthilkumar B. Hepatoprotective potential of *Andrographis paniculata* aqueous leaf extract on ethanol induced liver toxicity in albino rats. *Journal of Applied Pharmaceutical Science*. 2011;1(6):24–28.
- [29] Asha T, Londonkar R, Nayaka H. Cytotoxicity and hepatoprotective attributes of methanolic extract of *Rumex vesicarius* L. *Biological Research*. 2015;48(1):1–9.
- [30] Kanchana G, Jayapriya K. Antioxidant effect of Livomap, A polyherbal formulation on ethanol induced hepatotoxicity in albino Wistar rats. *Journal of Applied Pharmaceutical Science*. 2013;3(3):52–56.
- [31] Babbar N, Oberoi H, Uppal D, Patil R. Oberoi HS, Uppal DS, Patil RT. Total phenolic content and antioxidant capacity of extracts obtained from six important fruit residues. *Food Research International*. 2011;44(1):391–396.
- [32] Kamali M, Khosroyar S, Kamali H, Ahmadzadeh ST, Mohammadi A. Phytochemical screening and evaluation of antioxidant activities of *Dracocephalum kotschyi* and determination of its luteolin content. *Avicenna Journal of Phytomedicine*. 2016;6(4):425–433.
- [33] Santhosh S, Sini K, Anandan R, Mathew P. Hepatoprotective activity of chitosan against isoniazid and rifampicin induced toxicity in experimental rats. *European Journal of Pharmacology*. 2007;57(2):69–73.
- [34] Kamel H, Azza H, Rehman A, Walaa M, Amira H. Protective effect of some antioxidants against CCl4 induced toxicity in liver cells from BRL3A cell line. *Journal of American Science*. 2010;6(10):992–1003.
- [35] Bhagyashree R, Sanjeevani H, Ujwala R. In vitro protection of hepatocytes by *Alocasia macrorrhiza* leaf juice against CCl4 and tylenol mediated hepatic injury. *International Journal of Pharmaceutical Applications*. 2011;2(2):122–128.